



University of Kentucky  
**UKnowledge**

---

International Grassland Congress Proceedings

XXI International Grassland Congress / VIII  
International Rangeland Congress

---

## Forage Potential of Legumes Native to Sub-Tropical Southeastern USA

Kenneth H. Quesenberry  
*University of Florida*

J. M. Dampier  
*University of Florida*

E. Valencia  
*University of Puerto Rico, Mayaguez*

Follow this and additional works at: <https://uknowledge.uky.edu/igc>

 Part of the [Plant Sciences Commons](#), and the [Soil Science Commons](#)

This document is available at <https://uknowledge.uky.edu/igc/21/13-1/23>

The XXI International Grassland Congress / VIII International Rangeland Congress took place in Hohhot, China from June 29 through July 5, 2008.

Proceedings edited by Organizing Committee of 2008 IGC/IRC Conference

Published by Guangdong People's Publishing House

---

This Event is brought to you for free and open access by the Plant and Soil Sciences at UKnowledge. It has been accepted for inclusion in International Grassland Congress Proceedings by an authorized administrator of UKnowledge. For more information, please contact [UKnowledge@lsv.uky.edu](mailto:UKnowledge@lsv.uky.edu).

## Forage potential of legumes native to sub-tropical southeastern USA

K.H. Quesenberry<sup>1\*</sup>, J.M. Dampier<sup>1</sup>, and E. Valencia<sup>2</sup>

<sup>1</sup>P.O. Box 110500, University of Florida, Gainesville, FL 32611; <sup>2</sup>University of Puerto Rico, Mayaguez.

\* Corresponding author (clover@ufl.edu)

**Key words** : native legumes ,centrosema ,crotalaria ,desmodium ,galactia ,Indigofera ,Tephrosia ,Trifolium

**Introduction** The subtropical part of the southeastern USA is one of the most difficult areas of the world for forage development . Climatic conditions range from tropical summer temperatures with high humidity to short periods of sub-freezing winter temperatures . Many forage legumes adapted to more temperate area have been evaluated for use in this region , but fail due to lack of summer persistence and successful pasture applications have been very limited . Likewise forage legume species from more tropical areas of South America and SE Asia often experience winter-kill and have found limited use . Additionally , root-knot nematodes ( *Meloidogyne* spp . ) ( RKN ) may dramatically limit legume persistence in lighter textured soils of this region . Research with the seven *Trifolium* spp . native to this region ( *T. bejariense* Moric . , *T. calcaricum* J . L . Collins & Wieboldt , *T. carolinianum* Michx . , *T. polymorphum* Poir . , *T. reflexum* L . , *T. stoloniferum* Muhl . ex Eaton , and *T. virginicum* Small ) , has shown that they generally have higher levels of RKN tolerance than introduced *Trifolium* species ( Quesenberry , et al . , 1997 ) . We hypothesize that evaluation of legume species native to the region and similar subtropical climatic zones will identify species with higher success potential .

**Materials and methods** One accession each of *Centrosema virginianum* ( L . ) Benth . , *Chamaecrista nictitans* ( L . ) Moench . , *Crotalaria rotundifolia* J . F . Gmel . . , *Desmodium incanum* DC . , *Desmodium lineatum* DC . , *Desmodium paniculatum* ( L . ) DC . , *Galactia regularis* ( L . ) Britton et al . . , *Indigofera spicata* Forssk . , *Tephrosia spicata* ( Walter ) Torr . & A . Gray and *Trifolium carolinianum* Michx . was evaluated for field performance and response to RKN . All germplasm was collected from native stands in the vicinity of Gainesville , FL , USA . Field evaluations consisted of spaced plants transplanted into herbicide suppressed bahiagrass ( *Paspalum notatum* Flugge ) sod . Multiple accessions of *D. incanum* obtained from the USDA NPGS were evaluated in the field . Evaluation of RKN response was conducted using previously published ( Quesenberry et al . , 1997 ) methods for greenhouse screening . Legumes were evaluated for response to *Meloidogyne arenaria* race 1 , *M. incognita* race 4 , or *M. javanica* . Additionally legumes were evaluated for tannin levels of field grown forage in both Florida and Puerto Rico .

**Results** There were differences (  $P < 0.05$  ) among legume species for root gall and egg mass scores in response to each of the three RKN species . In general , the level of root galling and egg mass production was lower than that observed on introduced forage legumes in previous research ( Kouame et al . , 1987 ) , although the accessions of *C. nictitans* , *D. lineatum* , and *T. spicata* would be classified as moderately to highly susceptible ( root gall scores  $\geq 3.0$  ) to all three nematodes . *Desmodium incanum* showed a near immune response to all three RKN species with no gall or egg masses observed . Similar to previously reported results *T. carolinianum* ( the only clover species native to this region ) showed highly resistant responses to these RKN species .

Field evaluations showed superior persistence for *D. incanum* which spread readily into an established bahiagrass sod . Although there was variability among accessions for productivity , this species in general showed good persistence over three years in the grass sod . However , a later planting of selected accessions in a bare soil became infected with an unidentified potyvirus and did not persist . Testing was conducted for BYMV , SMV and WMV , but not BCMV or DMV . *Centrosema virginiana* also showed good persistence in bahiagrass sod .

Tannin levels varied among accessions of *D. incanum* and among the other species evaluated . Measured levels were similar to accessions of *D. heterocarpon* that have been reported by other researchers under tropical conditions . The species of *Centrosema* , *Tephrosia* , *Galactia* , *Crotalaria* , and *Lotononis* showed lower levels of tannins than the *Desmodium* species .

**Conclusions** Domestication of plant species for cultivated agriculture has occurred over many centuries . Identification of legume native to a region may identify species having general ecologically adaptive traits and specific resistances to pest prevalent in the region . This research showed that legumes native to the SE USA in general did have high levels of RKN resistance and some of them were persistent when grown in a bahiagrass sod . However , other pest such as plant viruses may limit their usefulness . One must also be aware of factors such as tannins that may be associated with pest resistance , but may have an anti-quality attribute in forage-livestock systems .

### References

- Quesenberry , K . H . , N . M . Call \* , D . E . Moon \* , and R . A . Dunn . 1997 . Response of native eastern North American *Trifolium* spp . to root-knot nematodes . *Crop Sci.* 37 :270-274 .  
Kouame \* , C . N . , K . H . Quesenberry , and R . A . Dunn . 1997 . Response to root-knot nematodes of a germplasm collection of red clover and related species . *Genet . Resour . & Crop Evol.* 44 :439-445 .